

Privacy Policy





S NCBI		PubyMea				of Medicine NLM			
Entrez	PubMed	Nucleotide	Protein	Genome	Structure	ОМІМ	PMC	Journals	В
Search	PubMed	for				Go	Clear		
		Limits	Previe	ew/Index	History	Cli	pboard	De	tails
About Enti	rez	Display	tract	Sho	w: 20 🔽	Sort	∵ Seno	to: Text	
Text Version		□1: Res Vet	Sci. 1981 N	May;30(3):2	61-5.			Related Arti	icles, I
Entrez PubMed Overview Help FAQ Tutorial New/Noteworthy E-Utilities		Location and secretion of gastric intrinsic factor in the sheep.							
		McKay EJ, McLeay LM.							
PubMed Services Journals Database MeSH Database Single Citation Matcher Batch Citation Matcher Clinical Queries LinkOut Cubby		The site of production and secretion of intrinsic factor (IF) in the sheep has be studied using a human auto-antibody directed against IF. Immunofluorescent studies indicated the abomasal parietal cell was the source of IF in the sheep. Concentrations of IF in pure gastric secretion of sheep were relatively stable to 4 iu/ml and it was estimated that the total abomasal output of IF was 10,00 to 23,500 iu per 24 hours.							
Related Resources Order Documents		PMID:	7020016 [P	ubMed - inc	lexed for Ml	EDLINE]			
NLM Gate TOXNET Consumer Clinical Al- ClinicalTri PubMed C	way ⁻ Health erts als.gov	Display Abs	stract	Sho	w: 20 ▼	Sort j	Sen	d to. Text	ţ

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

Jan 29 2004 15: